

Osteoarthritis and Cartilage



KOOS pain as a marker for significant knee pain two and six years after primary ACL reconstruction: a Multicenter Orthopaedic Outcomes Network (MOON) prospective longitudinal cohort study

D. Wasserstein †*, L.J. Huston ‡, S. Nwosu §, MOON Group^a, C.C. Kaeding ¶, R.D. Parker #, R.W. Wright ††, J.T. Andrich †††, R.G. Marx §§, A. Amendola |||, B.R. Wolf |||, E.C. McCarty ¶¶, M. Wolcott ¶¶, W.R. Dunn ##

¶ Ohio State University, Columbus, OH, USA

Cleveland Clinic Foundation, USA

†† Washington University in St. Louis, USA

‡‡ Cleveland Clinic Foundation, Cleveland, OH, USA

§§ Hospital for Special Surgery, New York, NY, USA

||| University of Iowa, Iowa City, IA, USA

¶¶ University of Colorado, Denver, CO, USA

University of Wisconsin, Madison, WI, USA

K.P. Spindler ||

† University of Toronto Sunnybrook Health Sciences Centre, 2075 Bayview Avenue, MG301, Toronto, Ontario M4N 3M5, Canada

‡ Vanderbilt Sports Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

§ Department of Biostatistics, Vanderbilt University Medical Center, Nashville, TN, USA

|| Cleveland Clinic Sports Health, Cleveland Clinic, Cleveland, OH, USA

ARTICLE INFO

Article history:

Received 22 July 2014

Accepted 26 May 2015

Keywords:

Symptomatic osteoarthritis

Anterior cruciate ligament reconstruction

Knee Injury and Osteoarthritis Outcome Score (KOOS)

Knee pain

SUMMARY

Objective: The prevalence of radiographic osteoarthritis (OA) after anterior cruciate ligament reconstruction (ACLR) approaches 50%, yet the prevalence of significant knee pain is unknown. We applied three different models of Knee injury and Osteoarthritis Outcome Score (KOOS) thresholds for significant knee pain to an ACLR cohort to identify prevalence and risk factors.

Design: Multicenter Orthopaedic Outcomes Network (MOON) prospective cohort patients with a unilateral primary ACLR and normal contralateral knee were assessed at 2 and 6 years. Independent variables included patient demographics, validated Patient Reported Outcomes (PRO; Marx activity score, KOOS), and surgical characteristics. Models included: (1) KOOS criteria for a painful knee = quality of life subscale <87.5 and ≥ 2 of: KOOS_{pain} <86.1, KOOS_{symptoms} <85.7, KOOS_{ADL} <86.8, or KOOS_{sports/rec} <85.0; (2) KOOS_{pain} subscale score ≤ 72 (≥ 2 standard deviations below population mean); (3) 10-point KOOS_{pain} drop from 2 to 6 years. Proportional odds models ($\alpha \leq 0.05$) were used.

Results: 1761 patients of median age 23 years, median body mass index (BMI) 24.8 kg/m² and 56% male met inclusion, with 87% (1530/1761) and 86% (1506/1761) follow-up at 2 and 6 years, respectively. At 6 years, $n = 592$ (39%), $n = 131$ (9%) and $n = 169$ (12%) met criteria for models #1 through #3, respectively. The most consistent and strongest independent risk factor at both time-points was

* Address correspondence and reprint requests to: D. Wasserstein, University of Toronto Sunnybrook Health Sciences Centre, 2075 Bayview Avenue, MG301, Toronto, Ontario M4N 3M5, Canada. Tel: 1-416-480-5798; Fax: 1-416-480-5886.

E-mail addresses: E-mail addresses: david.wasserstein@sunnybrook.ca, david.wasserstein@utoronto.ca (D. Wasserstein), laura.huston@vanderbilt.edu (L.J. Huston), sam.nwosu@vanderbilt.edu (S. Nwosu), christopher.kaeding@osumc.edu (C.C. Kaeding), parkerr@ccf.edu (R.D. Parker), rwwright1@aol.com (R.W. Wright), andrisj@ccf.org (J.T. Andrich), marxr@hss.edu (R.G. Marx), ned-amendola@uiowa.edu (A. Amendola), brian-wolf@uiowa.edu (B.R. Wolf), eric.mccarty@ucdenver.edu (E.C. McCarty), michelle.wolcott@ucdenver.edu (M. Wolcott), dunn@ortho.wisc.edu (W.R. Dunn), spindlk@ccf.org (K.P. Spindler).

^a c/o laura.huston@vanderbilt.edu.

subsequent ipsilateral knee surgery. Low 2-year Marx activity score increased the odds of a painful knee at 6 years.

Conclusions: Significant knee pain is prevalent after ACLR; with those who undergo subsequent ipsilateral surgery at greatest risk. The relationship between pain and structural OA warrants further study.

© 2015 Osteoarthritis Research Society International. Published by Elsevier Ltd. All rights reserved.

Introduction

Anterior Cruciate Ligament (ACL) reconstruction (ACLR) is the most effective and reproducible treatment for ACL injured patients who want to return to cutting and pivoting sports^{1–3}. More than half of the patients undergoing ACLR will have concomitant pathology, including injuries to the articular cartilage in more than 20%, lateral meniscal tears in up to 46% and medial meniscal tears in 38%⁴.

An ACL tear is a known risk factor for the development of osteoarthritis (OA)⁵. Intermediate and long-term follow-up of ACLR patients has demonstrated a high prevalence of radiographic findings consistent with post-traumatic OA^{6–8}. Which factors, including concomitant pathology, the original injury, surgical techniques, or other as yet unidentified factors, are most responsible for the development of radiographic changes is unknown. A systematic review of studies including patients 5- to 10-years after ACLR⁶, found radiographic joint space narrowing in 0–13% of patients with intact menisci, and 21–48% in those who had undergone either meniscectomy or repair. The meniscal status was also demonstrated to be important in a systematic review of non-reconstructed ACL injured patients⁷. Most studies, however, are limited by poor follow-up and significant heterogeneity in the classification systems utilized to describe radiographic OA.

Although the definitions can be challenging^{9,10}, a systematic review in 2011 demonstrated a relationship between structural OA and symptomatic OA among high quality studies¹¹. Studies using Osteoarthritis Initiative (OAI) data have yielded further insight. Oak *et al.*¹² found a correlation between joint space narrowing at study entry, and greater progression of narrowing over the course of the study, with worse Patient Reported Outcomes (PRO) at 4 years. Others have found weak correlations between PRO and magnetic resonance image (MRI) confirmation of joint space narrowing¹³, but these correlations were highest for the knee pain subscale of the Knee Injury and Osteoarthritis Outcome Score (KOOS).

A consensus expert panel developed a definition of patients with a symptomatic knee significant enough to seek medical attention. This definition, based on threshold levels of KOOS subscale scores¹⁴, was based on the long-term follow-up of patients who previously underwent isolated partial meniscectomy with intact cruciate ligaments. Other criteria for clinically significant knee pain that have been developed based on PRO, include the KOOS Minimal Clinically Important Difference (MCID) of 8–10 points¹⁵, and the Osteoarthritis Research Society International (OARSI) Standing Committee criteria for interventions of OA of the knee (“OARSI responder criteria”) of 20 points^{16,17}.

Given that many patients who undergo ACLR develop radiographic OA, the main objective of this study was to identify the prevalence of significant knee pain by PRO after ACLR, using published definitions and cut-offs for either symptomatic OA or clinically significant knee pain. The second objective was to identify risk factors for developing a painful knee from patient, injury, and surgical characteristics 6 years following an ACL reconstruction.

Methods

Study design

Longitudinal prospective cohort (prognostic): The Multicenter Orthopaedic Outcomes Network (MOON) cohort¹⁸. MOON is a prospective, longitudinal, multicenter cohort study based in the United States, and designed to examine short and long-term prognosis after ACL reconstruction using validated patient-reported outcomes. MOON was also designed to generate hypotheses surrounding novel methods for improving outcomes after ACL injury.

Data sources

Participants completed a 13-page questionnaire providing patient demographics, a description of their injury, sports participation history, comorbidities and past medical history. Each participant also completed validated general and knee specific instruments, including the KOOS¹⁹ and the Marx activity rating scale²⁰. Contained within the KOOS is the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)²¹. All were completed within 2 weeks of the surgery date.

Surgeons completed a standardized questionnaire, which included detailed information regarding surgical technique, graft choice, and concomitant meniscal and articular cartilage pathology and treatment. The inter-rater reliability of grading systems for articular cartilage (modified Outerbridge) and meniscal lesions were previously validated among participating surgeons and found to be high^{22,23}. Meniscal pathology was classified by size, location, partial vs complete tears and treatment (not treated, repaired, resection and extent of resection).

Cohort design

All patients ($n = 2222$) who had undergone a unilateral primary ACLR at a participating MOON institution (Vanderbilt University, The Ohio State University, Washington University at St. Louis, University of Iowa, the Cleveland Clinic, and the Hospital for Special Surgery) between 2002 and 2005 were eligible for inclusion into this study. All patients provided informed consent from their respective institution. A prior exclusion criteria included previous contralateral ACL reconstruction, simultaneous bilateral ACL reconstruction, ACL repair, or a revision ACL reconstruction as the index (enrollment) event. ACL revision patients report worse PRO than primary ACL reconstruction patients^{4,24–26}, and so were excluded. ACL repair is an atypical treatment and was excluded. Patients with a contralateral ACL reconstruction prior to initial enrollment into the MOON cohort, or performed concurrently, were excluded on the basis that this study's objective included understanding how a subsequent contralateral reconstruction would influence PRO for significant knee pain. No patients were excluded from analysis due to incomplete baseline data (all $n = 2222$ completed baseline PRO).

Outcomes – definitions of a ‘painful or symptomatic knee’

We utilized previously published definitions of KOOS thresholds for a symptomatic knee as described in the introduction. We built three models, as follows:

- Model #1.** The primary model was defined as the operational definition of Englund *et al.*¹⁴ (“Englund model”) to distinguish patients with sufficient knee symptoms to seek medical care. The Englund model is defined as having a KOOS knee-related quality of life (QoL) subscale ≤ 87.5 and two or more of the other subscales: KOOS pain ≤ 86.1 , KOOS symptoms ≤ 85.7 , KOOS activities of daily living (ADL) ≤ 86.8 , or KOOS sports and recreation (“sport/rec”) ≤ 85.0 .
- Model #2.** The KOOS knee pain subscale has been shown to have the highest correlation with structural OA changes¹³, and is a direct measure of knee pain. Therefore, we defined a secondary model for significant knee pain as a KOOS pain subscale two standard deviations lower than the reported normal mean value in athletic populations with a history of (any) knee ligament injury. This value was 92.3 ± 10.0 ²⁴, which translated into a cut-off score of ≤ 72 points (“KOOS pain ≤ 72 model”). This definition also qualified as a 20-point change, consistent with OARSI responder criteria for effective interventions in OA¹⁶.
- Model #3.** The reported MCID for the KOOS pain subscale is 6.1 points in athletes after ACL reconstruction²⁷, to between 8 and 10 points for patients with OA^{15,28}. To utilize a more conservative estimate of the MCID, we selected a drop of 10 points in the KOOS pain subscale from 2 years to 6 years follow-up as an additional secondary definition of patients with a painful knee after ACL reconstruction (“KOOS pain MCID model”). This model attempted to identify patients who had a clinically significant worsening of knee pain.

Model variables

Variables included all those from the original MOON cohort. They included patient demographics (age, sex, body mass index [BMI], smoking status, education level, main sport played at the time of injury, enrollment year), validated PRO (KOOS, WOMAC, Marx activity), surgical characteristics (graft type, meniscal pathology/treatment, articular cartilage pathology), and incidence of subsequent surgery on either knee (Table I). The Marx score is a measure of the frequency and intensity of cutting and pivoting sports. The inclusion of variables in our models was based on substantive knowledge of the clinical or epidemiological association between them and PRO after ACL reconstruction surgery. These relationships have been established by our own work with this cohort^{4,7,18,24} and have been derived from literature review¹⁸. They extend to include baseline PRO scores, patient demographic factors and surgical variables.

Statistical analysis

To describe our patient sample, we summarized continuous variables as percentiles (i.e., 25th, 50th, and 75th) with their mean and standard deviation, and categorical variables with frequencies and percentages. Multivariable regression analyses were constructed to examine which baseline risk factors were independently associated with each outcome variable. An *a priori* determined list of variables to be included in all models were given by: age, gender, BMI, smoking status, education level, main sport played the last 2 years, baseline KOOS, WOMAC, and Marx activity levels, graft type, previous meniscal pathology, current meniscal

pathology/treatment, previous articular cartilage pathology, current articular cartilage pathology, subsequent surgery on the ipsilateral and contralateral knee, and enrollment year. We assumed independence of all covariates because we compared between subjects and not within, and when fitting the multivariable regression models, we measured each covariate’s independent adjusted association with the outcome. For binary outcome variables a multivariable logistic regression model was fit to the data, parameter estimates were exponentiated to obtain odds ratios (OR) and 95% confidence intervals (CI), based on a dichotomous outcome (yes/no). We did not assume a linear relationship between continuous covariates (independent variables) and each outcome in order to avoid underestimating the true relationship, instead utilized a restricted cubic regression splines technique that assumes smooth relationships (i.e., they are linearly related to the log odds). To avoid case-wise deletion of records with missing covariates, we employed multiple imputation *via* predictive mean matching. All model assumptions (as listed above) were met. Statistical analysis was performed using open source R statistical software (www.r-project.org; Version 3.0.3).

Post hoc analysis

Preliminary findings demonstrated that a low Marx activity score at 2 years increased the odds of reporting a painful knee in both the Englund and KOOS pain ≤ 72 models. Therefore, in order to further understand the interaction of pain and activity, we performed a *post hoc* analysis to identify the proportion of patients reporting a high level of sport/activity-related knee pain, and to understand which factors modified that outcome. This model utilized responses from a 5-point Likert question on the International Knee Documentation Committee (IKDC): “What is the highest level of activity that you can perform without significant knee pain?” Patients were classified based on their answer to this question as high activity tolerance (“very strenuous activities” or “strenuous activities” or “moderate activities”) or low activity tolerance (“light activities” or “no described activities”). Models were built for this outcome (**Model #4**: “Activity tolerance model”) at 6 years based on the response to the question at 2 years.

After determining that subsequent ipsilateral surgery was a risk factor, we performed a second post-hoc analysis to identify the number of patients who underwent a second surgery within 1 and 3 months prior to the 2- and 6-year time-points. This was performed due to concern that recent surgery may be the cause of higher reported pain. Furthermore, we re-analyzed each of the four models after excluding the patients with surgery within 3 months of the 2- and 6-year time-points.

Results

Figure 1 illustrates the cohort inclusion/exclusion criteria. There were 1761 subjects who fit the inclusion criteria and were included in this analysis. The median age of our cohort was 23 years, median BMI 24.8 kg/m² and the cohort was 56% male. Patient follow-up was obtained on 87% (1530/1761) and 86% (1506/1761) at 2 and 6 years, respectively. The proportion of patients who met each of the three model criteria was calculated (see Table III). At 2 years, $n = 46$ patients fit both the Englund and KOOS pain ≤ 72 points models, out of a total $n = 141$ possible patients (32.6%). At 6 years, $n = 67$ patients fit all four models, out of a total $n = 131$ possible (51%). Full baseline demographics are supplied in Table II alongside the list of model variables and levels.

Table III depicts the significant independent risk factors identified in each model. Subsequent ipsilateral surgery was the most consistent and strongest predictor of increased symptoms at both 2

Table 1
List of modeling variables

	Category	Levels
Baseline outcome scores	KOOS (5 subscales); WOMAC (pain, stiffness subscales)	Continuous
Patient demographics	Age (years)	Continuous
	Gender	Male, female
	BMI	Continuous
	Smoking status	Never, quit, current
	Education level (years)	1–16 (continuous)
	Baseline activity level (Marx)	Continuous
Surgical variables	Main sport played last 2 yrs	Basketball, football, soccer, other, none
	Graft type	Autograft (BTB), autograft (soft tissue), allograft (BTB), allograft (soft tissue)
	Meniscal pathology	
	* Previous	No, yes
	* Medial	Normal, no tx for tear, repair, excised, other
	* Lateral	Normal, no tx for tear, repair, excised, other
	Articular cartilage pathology	
	* Previous	No, yes
	* Medial femoral condyle (MFC)	Normal/grade 1, grade 2, grade 3, grade 4
	* Lateral femoral condyle (LFC)	Normal/grade 1, grade 2, grade 3, grade 4
* Medial tibial plateau (MTP)	Normal/grade 1, grade 2, grades 3/4	
* Lateral tibial plateau (LTP)	Normal/grade 1, grade 2, grades 3/4	
* Patella	Normal/grade 1, grade 2, grades 3/4	
* Trochlea	Normal/grade 1, grade 2, grades 3/4	
Miscellaneous variables	Year of surgery (enrollment)	2002, 2003, 2004, 2005
	Subsequent ipsilateral surgery	No, arthroscopic procedure, revision ACL reconstruction, total knee arthroplasty (TKA)
	Subsequent contralateral surgery	No, arthroscopic procedure, ACL reconstruction, total knee arthroplasty (TKA)

and 6 years post-ACL reconstruction (broken down by type in Fig. 2). Subsequent surgeries were common, occurring at a rate of 16% (239/1530) at 2 years and 21.5% (324/1506) at 6 years. The majority of subsequent surgeries other than total knee replacement took place more than a year prior to the 6 year outcome measurement. The mean time to revision ACLR was 2.4 ± 1.9 years, total

knee replacement 5.3 ± 2.2 years and other arthroscopic surgical procedures 2.1 ± 1.9 years. Revision ACLR was the single most common subsequent procedure. The vast majority of subsequent procedures took place remote from sampled time-points: only 1.3% (3/239) and 0% (0/324) of patients had a subsequent surgery within 1 month of filling out the KOOS forms at 2 years and 6 years,


**Fig. 1.** Flowchart showing the inclusion of participants in the study.

Table II
Baseline data for the included cohort and the patients lost to follow-up

Category	Variable (N)	Level	Overall cohort (n = 1761) n (%) or median (25th–75th)	Lost to follow-up @ 2 yrs (n = 231) n (%) or median (25th–75th)	Lost to follow-up @ 6 yrs (n = 255) n (%) or median (25th–75th)
Patient demographics	Sex	Male	980 (56%)	146 (63%)	169 (66%)
		Female	781 (44%)	85 (37%)	86 (34%)
	Age	Continuous	23 years (17–35)	22 years (17–30)	22 years (17–32)
		BMI	Continuous	24.8 kg/m ² (22.3–27.9)	25.4 kg/m ² (22.9–29.2)
	Smoking status	Current	167 (10%)	32 (15%)	44 (18%)
		Quit smoking (>6 months)	172 (10%)	15 (7%)	16 (6%)
		Never smoker	1354 (80%)	171 (78%)	186 (76%)
	Education level	Continuous	14.0 years (11.0–16.0)	13.0 years (11.0–16.0)	12.0 years (10.0–16.0)
		Main sport	Basketball	393 (23%)	61 (27%)
	Football		191 (11%)	38 (17%)	35 (14%)
	Soccer		230 (13%)	17 (7%)	20 (8%)
	Other		793 (46%)	99 (43%)	106 (42%)
	None		134 (8%)	14 (6%)	22 (9%)
PRO	Marx activity	Baseline	12 (8–16)	13 (9–16)	13 (8–16)
		2 years	9 (4–13)	N/A	8 (1–12)
	KOOS symptoms	Baseline	68 (57–82)	68 (50–79)	68 (50–82)
		KOOS pain	Baseline	75 (64–89)	69 (58–86)
	KOOS ADL	Baseline	88 (74–96)	83 (68–94)	82 (65–94)
		KOOS Sports & rec	Baseline	50 (30–75)	50 (25–75)
	KOOS QoL	Baseline	38 (25–50)	31 (19–50)	31 (19–44)
		WOMAC stiffness	Baseline	75 (62–88)	75 (50–88)
WOMAC pain	Baseline	90 (75–95)	85 (70–95)	85 (65–95)	
Surgical/Injury factors	Previous meniscal pathology	No	1632 (93%)	210 (91%)	230 (90%)
		Yes	129 (7%)	21 (9%)	25 (10%)
	Previous articular cartilage pathology	No	1739 (99%)	228 (99%)	249 (98%)
		Yes	22 (1%)	3 (1%)	6 (2%)
	Graft type	Allograft (BTB)	121 (7%)	16 (7%)	27 (11%)
		Allograft (soft tissue)	299 (17%)	39 (17%)	35 (14%)
		Autograft (BTB)	832 (47%)	116 (51%)	120 (47%)
		Autograft (soft tissue)	509 (29%)	60 (26%)	73 (29%)
	Medial meniscus treatment	Normal/none	1106 (63%)	149 (65%)	173 (68%)
		Tear/no treatment	94 (5%)	10 (4%)	10 (4%)
		Repair	229 (13%)	39 (17%)	35 (14%)
		Partial excision	317 (18%)	33 (14%)	37 (15%)
		Other	15 (1%)	0 (0%)	0 (0%)
	Lateral meniscus treatment	Normal/none	927 (53%)	118 (51%)	127 (50%)
		Tear/no treatment	197 (11%)	25 (11%)	25 (10%)
		Repair	128 (7%)	16 (7%)	22 (9%)
		Partial excision	497 (28%)	72 (31%)	80 (31%)
		Other	12 (1%)	0 (0%)	1 (<1%)
	Medial femoral condyle	Normal/grade 1	1381 (78%)	189 (82%)	202 (79%)
		Grade 2	225 (13%)	30 (13%)	31 (12%)
		Grade 3	117 (7%)	8 (3%)	15 (6%)
		Grade 4	38 (2%)	4 (2%)	7 (3%)
	Lateral femoral condyle	Normal/grade 1	1498 (85%)	197 (85%)	209 (82%)
		Grade 2	186 (11%)	24 (10%)	32 (13%)
		Grade 3	59 (3%)	10 (4%)	11 (4%)
		Grade 4	18 (1%)	0 (0%)	3 (1%)
	Medial tibial plateau	Normal/grade 1	1694 (96%)	228 (99%)	250 (99%)
		Grade 2	45 (3%)	1 (<1%)	1 (<1%)
		Grades 3–4	22 (1%)	0 (0%)	0 (0%)
	Lateral tibial plateau	Normal/grade 1	1611 (91%)	215 (94%)	230 (92%)
		Grade 2	122 (7%)	13 (6%)	20 (8%)

(continued on next page)

Table II (continued)

Category	Variable (N)	Level	Overall cohort (n = 1761) n (%) or median (25th–75th)	Lost to follow-up @ 2 yrs (n = 231) n (%) or median (25th–75th)	Lost to follow-up @ 6 yrs (n = 255) n (%) or median (25th–75th)
Patella		Grades 3–4	28 (2%)	0 (0%)	0 (0%)
		Normal/grade 1	1470 (83%)	199 (90%)	213 (88%)
		Grade 2	181 (10%)	21 (10%)	29 (12%)
Trochlea		Grades 3–4	110 (6%)	0 (0%)	0 (0%)
		Normal/grade 1	1641 (93%)	219 (96%)	238 (96%)
		Grade 2	82 (5%)	10 (4%)	11 (4%)
		Grades 3–4	38 (2%)	0 (0%)	0 (0%)

respectively. Furthermore, only 3.8% (9/239) and 1.2% (4/324) of patients had a subsequent surgery within 3 months of filling out the KOOS forms at 2 years and 6 years, respectively. When all four models were re-run with patients who had undergone subsequent surgery within 3 months removed, no changes were noted in the significance or magnitude of any statistically significant risk factors. One risk factor in model #4 which previously approached significance then became significant (current vs never smoker: OR 1.82 (1.02, 3.27); $P = 0.043$).

Other independent risk factors that were found to be significant (although inconsistent) of increased symptoms at 2 and/or 6 years post-ACL reconstruction included higher BMI, smokers, less years of education, lower baseline KOOS ADL and higher baseline KOOS sports/rec subscale scores, and lower 2-year Marx activity levels (for predicting the 6-year models).

Potential prognostic factors that did not alter the risk of reporting a painful knee or having significant activity-related pain included age, pre-operative/baseline activity level, pre-operative WOMAC (pain, stiffness) or KOOS (symptoms, pain, and QoL) baseline scores, graft type, medial meniscal pathology/treatment, and subsequent contralateral knee surgery. The grade of chondral damage at initial arthroscopy was an inconsistent predictor in the patellofemoral, medial and lateral compartments. In general, when chondral damage influenced the odds of reporting either a painful knee or significant activity-related pain, the tendency was for the effect to be driven by grade 3/4 change.

Discussion

The prevalence of significant patient-reported knee pain 6 years after ACLR was high, including 39% by the Englund definition, 9% for KOOS pain score ≤ 72 (drop ≥ 20 points) and 12% for KOOS pain MCID definition (drop ≥ 10 points). A similar proportion of patients (11%) reported significant activity-related knee pain at 6 years. This study is the first to apply these definitions to characterize this patient population and represents an important first step in identifying at-risk patients for the development of significant knee pain after ACLR.

The most consistent risk factor across all definitions of significant knee pain also carried the largest impact – subsequent ipsilateral knee surgery. We utilized interactions of age and subsequent surgery in our statistical modeling because of our previous findings⁴ that demonstrated younger age increased the risk of subsequent surgery at 2 and 6 year follow-up. This limited the degrees of freedom we could use to identify which of the subsequent procedures had the most influence. At 6 years, ipsilateral re-operation was dominated by revision ACLR, further meniscus/articular cartilage surgery and surgical interventions for stiffness. Revision ACLR has been associated with worse PRO^{4,24–26}, and subsequent meniscal or articular cartilage surgery is a recognized risk for radiographic OA changes in ACL reconstructed patients⁵. The identification of subsequent surgery as a risk factor for reporting a painful knee was also a robust enough finding that it held even with the removal of patients who had surgery within 3 months of the 2- and 6-year time-points from statistical analysis. That contralateral knee surgery did not increase the odds of reporting a painful knee, places further importance on subsequent surgeries as a marker for additional trauma or joint degeneration as a driver of poor outcomes. Better resolution of the type of procedure in subsequent investigations will be helpful, as some are potentially preventable through improved surgical technique, timing of surgery, or rehabilitation.

Many ACLR patients exhibited activity-related knee pain in follow-up. We assessed this using model #4, and determined that 11% of patients met these criteria. This included approximately half

Table III
Summary of Independent Variables [reported as OR (95% CI); bolded and *P*-value included when significant]

Category	Variable comparison	Variable (worse outcome, if significant)	Model 1 (Englund)		Model 2 (KOOS Pain ≤ 72)		Model 3 (KOOS Pain MCID)	Model 4 (Activity tolerance)	
			2 years	6 years	2 years	6 years	6 years	6 years	
Number of patients who satisfied criteria for each model			655/1527 (43%)	592/1505 (39%)	141/1528 (9%)	131/1504 (9%)	169/1394 (12%)	159/1495 (11%)	
Baseline outcome scores	KOOS	Symptoms	0.90 (0.69, 1.18)	0.83 (0.69, 1.09)	0.78 (0.45, 1.36)	0.81 (0.45, 1.49)	0.91 (0.59, 1.40)	1.27 (0.78, 2.07)	
		Pain	0.66 (0.40, 1.07)	0.78 (0.46, 1.32)	0.48 (0.17, 1.36)	0.87 (0.32, 2.37)	1.45 (0.66, 3.18)	1.00 (0.40, 2.51)	
		ADL	0.53 (0.36, 0.79) <i>P</i> < 0.001	0.67* (0.45, 0.99) <i>P</i> = 0.117	0.78 (0.37, 1.62)	0.89 (0.42, 1.89)	1.13 (0.61, 2.07)	0.67 (0.32, 1.41)	
		Sports/Rec	1.04 (0.82, 1.32)	1.01 (0.77, 1.33)	1.63 (1.13, 2.34) <i>P</i> = 0.016	0.78 (0.53, 1.16)	0.90 (0.63, 1.30)	0.88 (0.60, 1.29)	
Patient characteristics	WOMAC	QoL	0.90 (0.76, 1.06)	0.97 (0.82, 1.15)	0.96 (0.73, 1.27)	1.00 (0.73, 1.36)	0.82 (0.64, 1.04)	0.79 (0.60, 1.04)	
		Pain	1.18 (0.77, 1.81)	1.09 (0.69, 1.72)	0.68 (0.31, 1.47)	0.46 (0.20, 1.05)	0.63 (0.29, 1.35)	1.03 (0.50, 2.13)	
		Stiffness	0.92 (0.74, 1.15)	0.92 (0.72, 1.16)	1.06 (0.70, 1.59)	1.57 (1.00, 2.47)	1.07 (0.73, 1.58)	0.92 (0.62, 1.36)	
Patient characteristics	Age (yrs)	Age	0.91 (0.55, 1.51)	0.76 (0.42, 1.37)	1.65 (0.73, 3.71)	1.21 (0.48, 3.05)	0.96 (0.42, 2.17)	1.43 (0.62, 3.28)	
	Gender	Females: Males	Males	1.10 (0.86, 1.40)	1.00 (0.75, 1.33)	0.85 (0.54, 1.34)	0.90 (0.54, 1.51)	0.67 (0.47, 0.96) <i>P</i> = 0.028	1.46 (0.89, 2.39)
	BMI	BMI	Higher BMI	1.24 (1.01, 1.53) <i>P</i> = 0.004	1.28* (1.01, 1.61) <i>P</i> = 0.071	1.52 (1.04, 2.20) <i>P</i> = 0.003	1.30 (0.84, 2.02)	1.00 (0.71, 1.39)	1.27 (0.89, 1.83)
	Smoking status	Current: Never	Current (compared to never)	1.45 (0.96, 2.19)	1.22 (0.81, 1.85)	1.59 (0.92, 2.75)	2.83 (1.46, 5.49) <i>P</i> = 0.002	1.13 (0.62, 2.08)	1.78 (1.00, 3.18)
		Quit: Never	Quitting (compared to never)	1.66 (1.12, 2.45) <i>P</i> = 0.011	0.84 (0.54, 1.32)	0.96 (0.50, 1.83)	1.96 (1.02, 3.75) <i>P</i> = 0.042	0.59 (0.32, 1.12)	1.03 (0.57, 1.89)
	Education (years)	Years of education	Less education years	1.04 (0.77, 1.41)	0.80 (0.56, 1.14)	0.63 (0.38, 1.03)	0.43 (0.23, 0.79) <i>P</i> = 0.022	0.72 (0.44, 1.16)	0.61* (0.39, 0.98) <i>P</i> = 0.126
	Baseline activity level (Marx)	Marx activity score		1.31 (0.92, 1.87)	1.16 (0.80, 1.69)	1.38 (0.72, 2.65)	1.39 (0.66, 2.92)	0.93 (0.53, 1.63)	1.26 (0.60, 2.67)
2 year activity level (Marx)	Marx activity score	Low score (low activity)	–	0.62 (0.48, 0.82) <i>P</i> = 0.001	–	0.53 (0.33, 0.86) <i>P</i> = 0.032	0.91 (0.64, 1.31)	0.41 (0.26, 0.63) <i>P</i> < 0.001	
Surgical factors	Previous meniscal pathology	Yes: No	No (compared to 'yes')	1.48 (0.91, 2.39)	1.23 (0.76, 1.98)	1.11 (0.57, 2.16)	1.13 (0.51, 2.52)	0.43 (0.19, 0.97) <i>P</i> = 0.041	1.78 (0.90, 3.51)
	Previous articular cartilage pathology	Yes: No		3.98 (0.95, 16.73)	0.70 (0.20, 2.50)	2.50 (0.70, 8.86)	1.79 (0.46, 6.93)	0.53 (0.08, 3.55)	3.47 (0.58, 20.63)
	Current meniscal pathology								
	* Medial	Repair: Normal		1.34 (0.94, 1.90)	1.11 (0.79, 1.55)	1.53 (0.90, 2.60)	1.61 (0.91, 2.86)	0.96 (0.59, 1.59)	1.21 (0.65, 2.25)
	* Lateral	No tear for treatment: Normal	Normal (compared to no tx for tear)	0.59 (0.39, 0.89) <i>P</i> = 0.012	0.74 (0.50, 1.10)	0.97 (0.50, 1.89)	0.62 (0.28, 1.40)	0.93 (0.52, 1.67)	1.22 (0.65, 2.30)
	Current Articular cartilage pathology								
	* Medial femoral condyle (MFC)	Grade 4: Normal/grade 1	Grade 4 (compared with normal/grade 1)	1.62 (0.69, 3.76)	1.43 (0.65, 3.14)	1.07 (0.35, 3.32)	1.28 (0.46, 3.61)	1.42 (0.50, 3.98)	2.67 (1.05, 6.80) <i>P</i> = 0.040
	* Lateral femoral condyle (LFC)	Grade 2: Normal/grade 1	Normal/grade 1 (compared to grade 2)	0.70 (0.47, 1.05)	0.73 (0.48, 1.09)	0.75 (0.36, 1.57)	0.34 (0.14, 0.82) <i>P</i> = 0.016	0.83 (0.46, 1.49)	1.01 (0.53, 1.92)
	Grade 3: Normal/grade 1	Grade 3 (compared with normal/grade 1)	1.78 (0.85, 3.73)	2.58 (1.21, 5.50) <i>P</i> = 0.014	1.40 (0.54, 3.62)	1.37 (0.44, 4.29)	1.89 (0.83, 4.32)	1.56 (0.47, 5.19)	
* Medial tibial plateau (MTP)	Grades 3/4: Normal/grade 1	Grades 3/4 (compared with normal/grade 1)	0.73 (0.24, 2.27)	2.98 (0.93, 9.50)	1.26 (0.35, 4.53)	3.20 (0.76, 13.48)	4.20 (1.33, 13.25) <i>P</i> = 0.015	0.86 (0.17, 4.25)	
* Lateral tibial plateau (LTP)	Grade 2: Normal/grade 1	Normal/grade 1 (compared to grade 2)	0.91 (0.57, 1.46)	0.46 (0.26, 0.82) <i>P</i> = 0.008	0.41 (0.15, 1.14)	1.24 (0.53, 2.93)	1.15 (0.60, 2.22)	0.56 (0.24, 1.35)	

(continued on next page)

Table III (continued)

Category	Variable comparison	Variable (worse outcome, if significant)	Model 1 (Englund)		Model 2 (KOOS Pain ≤ 72)		Model 3 (KOOS Pain MCID)		Model 4 (Activity tolerance)	
			2 years	6 years	2 years	6 years	6 years	6 years	6 years	6 years
* Patella	Grade 2: Normal/grade 1	Grade 2 (compared with normal/grade 1)	0.95 (0.63, 1.42)	1.65 (1.06, 2.58) P = 0.028	1.81 (0.96, 3.39)	1.59 (0.74, 3.43)	1.30 (0.66, 2.59)	3.06 (1.68, 5.56) P < 0.001		
	Grades 3/4: Normal/grade 1	Grades 3/4 (compared with normal/grade 1)	1.71 (1.04, 2.80) P = 0.033	1.55 (0.94, 2.55)	1.28 (0.60, 2.70)	2.29 (0.92, 5.68)	2.10 (1.09, 4.03) P = 0.026	2.14 (1.07, 4.30) P = 0.032		
* Trochlea	Grade 2: Normal/grade 1	Normal/grade 1 (compared to grade 2)	0.93 (0.52, 1.67)	0.96 (0.54, 1.73)	0.88 (0.37, 2.10)	0.44 (0.14, 1.37)	0.57 (0.22, 1.48)	0.31 (0.12, 0.82) P = 0.017		
Subsequent surgery	Ipsilateral knee	Yes: No	2.31 (1.35, 3.96) P < 0.001	2.66 (1.77, 4.01) P < 0.001	2.20 (1.07, 4.55) P < 0.001	3.41 (1.72, 6.75) P < 0.001	1.94* (1.11, 3.39) P = 0.060	3.03 (1.59, 5.79) P < 0.001		
	Contralateral knee	Yes: No	0.62 (0.27, 1.41)	0.92 (0.50, 1.68)	0.23 (0.03, 1.88)	0.90 (0.29, 2.76)	1.55 (0.54, 4.45)	0.47 (0.14, 1.64)		

Key:

Significant values are depicted in bold.

Gray shading indicates that outcome is counterintuitive to what one would think.

* Although OR did not cross 1, indicating significance, the P value did not reach 0.05 level of significance.

** After removal of patients with surgery ≤ 3 months before the outcome time-point, variable became significant (OR 1.82 (1.02, 3.27); P = 0.043).

For continuous variables, where the OR < 1.0 – variables are inversely related.

For categorical variables, where the OR > 1.0 – first variable listed is worse than second variable listed.

For categorical variables, where the OR < 1.0 – second variable listed is worse than first variable listed.

Subsequent ipsilateral procedures at 6 years




Fig. 2. Subsequent ipsilateral surgical procedures at 6 years (“Other arthroscopic” includes hardware removal, meniscal and articular cartilage surgery, infection, arthrolysis/manipulation).

who also met criteria for models #2 and #3 – both KOOS pain models. KOOS pain assesses both activity-related and non-activity related pain and contains the questions from the validated hip and knee OA tool – WOMAC¹⁵. Furthermore, we noted that a low Marx score at 2 years increased the odds of a patient meeting the Englund criteria (model #1), KOOS pain ≤ 72 criteria (model #2), and IKDC activity-related pain threshold (model #4). Whether simply being less active is a risk factor for reporting significant knee pain, or whether patients already developing significant knee pain become less active, is not known.

There is no consensus definition for symptomatic OA or significant knee pain using PRO. This is further complicated by the heterogeneity of diagnoses/definitions reported in the literature. While we found the prevalence of significant knee pain was high, it varied considerably based on our definitions. The Englund *et al.*¹⁴ criteria were the least stringent, but also the broadest including pain, symptoms and QoL reporting. The KOOS pain threshold in that model was 86 points, which corresponded to the 25th percentile of KOOS pain scores in the MOON cohort⁴. Accordingly, this model identified the most patients. Few cues are available from the literature for prevalence of pain based on the Englund model in similar patients, with only two small published studies. In a cohort of 84 female soccer players with an ACL injury treated with either rehabilitation or surgery²⁹, 75% met the Englund criteria at 15 years follow-up. In contrast, 51% met criteria for knee OA on radiographs, and 42% met both. In a purely non-operative cohort³⁰ of 67 ACL injured patients, the 15-year KOOS pain scores were all 85 points or greater.

The 20-point drop in KOOS pain score that we selected corresponded to the OARSI responder criteria¹⁶ and 2 standard deviations below the mean of KOOS scores of athletes with a history of knee ligament injury³¹. Even fewer comparative studies exist in the literature for this definition. Paradowski *et al.*¹⁷ applied the OARSI responder criteria, developed for use in OA interventional studies to identify therapies that produce significant knee pain reduction, to identify mild OA patients with significantly increased knee symptoms post-meniscectomy. Those with radiographic changes had a larger drop in KOOS pain score (11 points), and by 6 years they determined that 7% of patients had a ≥ 20 point KOOS pain drop. Another study of older, post-meniscectomy patients with intact ligaments demonstrated a mean baseline KOOS pain of 84 points³², but with high individual variation. Seven years later the same patients reported a further six point drop in KOOS pain on average which was worse in females and those with radiographic changes.

A 20-point KOOS pain drop that is two standard deviations below population norms³¹ should theoretically include only 2.5% of our cohort. In fact, however, the distribution of KOOS pain was skewed to the left at 6 years with more than 9% of patients having a score below this cut-off. This finding offers both clinical and statistical significance and reinforces the role of subsequent injury, joint degeneration, or concomitant pathology at the primary reconstruction in the identification of patients at-risk for high levels of self-reported knee pain. Furthermore, a significant proportion of these patients reported high levels of activity-related pain according to our IKDC model #4 definition.

Limitations

There are some challenges in comparing our cohort with previous studies that have attempted to develop and characterize the prevalence of significant knee pain. Prior studies have examined patients with a different primary surgical intervention – namely meniscectomy^{14,17,32,33}. The etiology of meniscal tears in those cohorts included both traumatic and atraumatic mechanisms, whereas our cohort had sustained a traumatic rupture of the ACL. Secondly, the meniscectomy cohorts have an older mean age than our cohort (mean age typically 46–56 years, compared to mean age <30 years at follow-up in our study). How a degenerative process and traumatic process modify the risk of developing knee OA is unknown.

Loss to follow-up in our study was 13% (2 year) or 14% (6 year). Although there is no consensus on the introduction of bias based on follow-up, most estimates suggest that <5% loss will have no effect, while >20% may pose serious threats to validity³⁴. Yet the direction and approximate magnitude of some covariates, such as socioeconomic markers, do not change with attrition approaching even 50%^{35,36}. MOON investigators go to considerable length to contact enrolled patients, including repeated mailings and phone calls. As noted in Table III, the proportion of males lost to follow-up was slightly higher (63% vs 56%), and some minor differences were seen for BMI and smoking status. We don't think this will have had a large effect on the study conclusions, as sex was not associated with outcome, smoking was inconsistently associated with only a couple outcomes and BMI was only a predictor in model #1.

Our study was not designed to identify the best definition for a symptomatic knee. Accordingly, we utilized various definitions, each with advantages and disadvantages as well as mixed support in the literature. The agreement between models was reasonable at 6 years after ACLR, as exemplified by identifying approximately half of the patients ($n = 67$) from the most stringent model (KOOS pain ≤ 72 ; $n = 131$) in the remaining models. The identification of which outcomes (pain, function, or ADL) remain most important to post-ACLR patients, and the establishment of cut-off scores using the

Patient Acceptable Symptomatic State (PASS) concept for ACLR will be important steps in further defining this subset of patients.

Finally, we did not have radiographs available in follow-up of these patients to correlate structural change with symptomatic findings, as has been done in smaller series post isolated meniscectomy^{17,29,32,37,38}. The interaction of structural changes with symptoms is an important area for future research. This is heightened by the discordance between our study and systematic reviews of post-ACLR reconstruction patients⁶ that suggests meniscal pathology at the time of injury/surgery moderates radiographic OA risk. It would appear that while meniscal loss initiates joint space changes, it may be a weaker mediator of symptoms compared to other factors we have identified such as chondral damage and subsequent injury/surgery. There is some support for this notion

based upon weak associations demonstrated between joint space narrowing and poor PRO in OAI cohort studies^{12,13}. A second explanation is that the follow-up in our study is not yet long enough for meniscal status at the time of surgery to have the same influence on PRO. Exploring these interactions in future work is of critical importance to define the patients truly at-risk for clinically relevant OA after ACL reconstruction.

Summary

Significant knee pain and symptoms is prevalent among 9–39% of first-time ACL reconstruction patients at 6 years. Patient-reported pain is affected to some degree by demographic factors and higher grades of concurrent cartilage damage at the index procedure, however, those who undergo second surgeries (e.g., revision, repeat arthroscopy) are at greatest risk. Whether this patient report of significant knee pain relates to structural arthritic changes requires further study.

Author contributions

Factor	Contributor
Study design/conception	DW, LJH, KPS
Acquisition of data	LJH
Analysis and interpretation of data	DW, LJH, SN, KPS
Drafting of manuscript	DW
Critical revision for intellectual content	DW, LJH, KPS
Final approval	DW, LJH, KPS
Provision of study materials or patients	AA, JTA, WRD, CCK, RM, ECM, RDP, KPS, MLW, BRW, RWW
Statistical expertise	SN
Obtaining of funding	KPS
Administrative, technical or logistic support	?
Collection and assembly of data	AA, JTA, WRD, LJH, CCK, RM, ECM, RDP, KPS, MLW, BRW, RWW

Responsible for integrity of the work:

- 1) Kurt P. Spindler (spindlk@ccf.org)
- 2) Laura J Huston (laura.huston@vanderbilt.edu)
- 3) David Wasserstein (david.wasserstein@mail.utoronto.ca)

Competing interest statement

David Wasserstein, Laura Huston, Samuel Nwosu, Kurt Spindler, and Warren Dunn report institutional funding by NIH/NIAMS grant #5R01 AR053684 (PI—Spindler) and #5K23 AR052392 (PI—Dunn). These same authors report institutional funding via Unrestricted Educational Gifts from Smith and Nephew Endoscopy and DonJoy Orthopaedics.

Robert Marx has received funds outside the submitted work from the Journal of Bone and Joint Surgery (American) as Associate Editor; royalties from Demos Health for The ACL Solution; and royalties from Springer for Revision ACL Reconstruction.

Annunziato Amendola has received personal funds outside the submitted work from Arthrex, Inc.; ArthroSurface, Inc.; and MTP Solutions.

Brian Wolf has received personal funds outside the submitted work from United Health Care for participation on the Scientific Advisory Board.

Rick W. Wright has received personal funds outside the submitted work for Board Membership on the American Board of Orthopaedic Surgery Board of Directors and from the American

Orthopaedic Association as Treasurer-Elect. Additionally, he has received personal funds outside the submitted work from the NIA/NIAMS in the form of a research grant, and book royalties from Wolters Kluwer Lippincott Williams & Wilkins.

Funding declaration

The MOON cohort has obtained internal and external funding from numerous peer reviewed and non-peer reviewed sources. These include National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases (Grant numbers: 5R01 AR053684; 5K23 AR052392), Center for Education and Research on Therapeutics/Agency of Health Research and Quality (Grant number: 5U18-HS016075). This project received partial support from the Orthopaedic Research and Education Foundation, DonJoy Orthopaedics, Smith and Nephew Endoscopy, Vanderbilt Sports Medicine Research Fund, and by CTSA award no. UL1TR000445 from the National Center for Advancing Translational Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent official views of the National Center for Advancing Translational Sciences or the National Institutes of Health. The funding sources have partially supported the development and maintenance of the database and provided funding for a percent of effort for some of the research personnel.

Acknowledgments

We thank all our funding sources for supporting this work. These include National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases (Grant numbers: 5R01 AR053684; 5K23 AR052392), Center for Education and Research on Therapeutics/Agency of Health Research and Quality (Grant number: 5U18-HS016075). This project received partial support from the Orthopaedic Research and Education Foundation, DonJoy Orthopaedics, Smith and Nephew Endoscopy, Vanderbilt Sports Medicine Research Fund, and by CTSA award no. UL1TR000445 from the National Center for Advancing Translational Sciences. The contents of this study are solely the responsibility of the authors and do not necessarily represent official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

We thank the research coordinators, analysts and support staff from the MOON sites, whose efforts related to regulatory, data collection, subject follow-up, data quality control, analyses, and manuscript preparation make this consortium possible.

We also thank all the subjects who generously enrolled and participated in this study.

References

1. Engebretsen L, Benum P, Fasting O, Molster A, Strand T. A prospective, randomized study of three surgical techniques for treatment of acute ruptures of the anterior cruciate ligament. *Am J Sports Med* 1990;18(6):585–90.
2. Grontvedt T, Engebretsen L, Bredland T. Arthroscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone grafts with and without augmentation. A prospective randomized study. *J Bone Jt Surg Br* 1996;78(5):817–22.
3. Spindler KP, Wright RW. Clinical practice. Anterior cruciate ligament tear. *N Engl J Med* 2008;359(20):2135–42.
4. Cox CL, Huston LJ, Dunn WR, Reinke EK, Nwosu SK, Parker RD, et al. Are articular cartilage lesions and meniscus tears predictive of IKDC, KOOS, and Marx activity level outcomes after ACL reconstruction? A 6-year MOON cohort study. *Am J Sports Med* 2014;42(5):1058–67.
5. Muthuri SG, McWilliams DF, Doherty M, Zhang W. History of knee injuries and knee osteoarthritis: a meta-analysis of observational studies. *Osteoarthr Cartil* 2011;19:1286–93.
6. Oiestad BE, Engebretsen L, Storheim K, Risberg MA. Knee osteoarthritis after anterior cruciate ligament injury. *Am J Sports Med* 2009;37(7):1434–43.
7. Magnussen RA, Mansour AA, Carey JL, Spindler KP. Meniscus status at anterior cruciate ligament reconstruction associated with radiographic signs of osteoarthritis at 5- to 10-year follow-up. *J Knee Surg* 2009;22(4):347–56.
8. Chalmers PN, Mall NA, Moric M, Sherman SL, Paletta GP, Cole BJ, et al. Does ACL reconstruction alter natural history? A systematic literature review of long-term outcomes. *J Bone Jt Surg Am* 2014;96(4):292–300.
9. Pereira D, Peleteiro B, Araujo J, Branco J, Santos RA, Ramos E. The effect of osteoarthritis definition on prevalence and incidence estimates: a systematic review. *Osteoarthr Cartil* 2011;19:1270–85.
10. Guermazi A, Roemer F, Felson D, Brandt KD. Motion for debate: osteoarthritis clinical trials have not identified efficacious therapies because traditional imaging outcome measures are inadequate. *Arthritis Rheum* 2013;65(11):2748–58.
11. Kinds MB, Welsing PM, Vignon EP, Bijlsma JW, Viergever MA, Marjinissen AC, et al. A systematic review of the association between radiographic and clinical osteoarthritis of hip and knee. *Osteoarthr Cartil* 2011;19(7):768–78.
12. Oak SR, Ghodadra A, Winalski CS, Miniaci A, Jones MH. Radiographic joint space width is correlated with 4-year clinical outcomes in patients with knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthr Cartil* 2013;21(9):1185–90.
13. Illingworth KD, Bitar YE, Seiwert K, Scaife SL, El-Amin S, Saleh KJ. Correlation of WOMAC and KOOS scores to tibiofemoral cartilage loss on plain radiography and 3 Tesla MRI: data from the osteoarthritis initiative. *Knee Surg Sports Traumatol Arthrosc* 2014;22(7):1649–58.
14. Englund M, Roos EM, Lohmander LS. Impact of type of meniscal tear on radiographic and symptomatic knee osteoarthritis. *Arthritis Rheum* 2003;48(8):2178–87.
15. Roos EM, Toksvig-Larsen S. Knee injury and Osteoarthritis Outcome Score (KOOS) – validation and comparison to the WOMAC in total knee replacement. *Health Qual Life Outcomes* 2003;1:17.
16. Dougados M, LeClaire P, van der Heijde D, Bloch DA, Bellamy N, Altman RD. Response criteria for clinical trials on osteoarthritis of the knee and hip. A report of the osteoarthritis research society international standing committee for clinical trials response criteria initiative. *Osteoarthr Cartil* 2000;8:395–403.
17. Paradowski PT, Englund M, Roos EM, Lohmander LS. Similar group mean scores, but large individual variations, in patient-relevant outcomes over 2 years in meniscectomized subjects with and without radiographic knee osteoarthritis. *Health Qual Life Outcomes* 2004;2:38.
18. Spindler KP, Parker RD, Andrich JT, Kaeding CC, Wright RW, Marx RG, et al. Prognosis and predictors of ACL reconstructions using the MOON cohort: a model for comparative effectiveness studies. *J Orthop Res* 2013;31(1):2–9.
19. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee injury and Osteoarthritis Outcome Score (KOOS): development of a self-administered outcome measure. *J Orthop Sports Phys Ther* 1998;28(2):88–96.
20. Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med* 2001;29(2):213–8.

21. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *J Rheumatol* 1988;15(12):1833–40.
22. Marx RG, Connor J, Lyman S, Amendola A, Andrish JT, Kaeding CK, et al. Multirater agreement of arthroscopic grading of knee articular cartilage. *Am J Sports Med* 2005;33(11):1654–7.
23. Dunn WR, Wolf BR, Amendola A, Andrish JT, Kaeding CK, Marx RG, et al. Multirater agreement of arthroscopic meniscal lesions. *Am J Sports Med* 2004;32(8):1937–40.
24. Wright R, Spindler K, Huston L, Amendola A, Andrish J, Brophy R, et al. Revision ACL reconstruction outcomes: MOON cohort. *J Knee Surg* 2011;24(4):289–94.
25. Wright RW, Gill CS, Chen L, Brophy RH, Matava MJ, Smith MV, et al. Outcome of revision anterior cruciate ligament reconstruction: a systematic review. *J Bone Jt Surg Am* 2012;94(6):531–6.
26. Lind M, Menhert F, Pedersen AB. Incidence and outcome after revision anterior cruciate ligament reconstruction: results from the Danish registry for knee ligament reconstructions. *Am J Sports Med* 2012;40(7):1551–7.
27. Salavati M, Akbhari B, Mohammadi F, Mazaheri M, Khorrami M. Knee injury and Osteoarthritis Outcome Score (KOOS); reliability and validity in competitive athletes after anterior cruciate ligament reconstruction. *Osteoarthr Cartil* 2011;19:406–10.
28. Roos EM, Lohmander LS. The Knee injury and Osteoarthritis Outcome Score (KOOS): from joint injury to osteoarthritis. *Health Qual Life Outcomes* 2003;1:64.
29. Lohmander L, Ostberg A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum* 2004;50(10):3145–52.
30. Kostogiannis I, Ageberg E, Neuman P, Dahlberg L, Friden T, Roos H. Activity level and subjective knee function 15 years after anterior cruciate ligament injury: a prospective, longitudinal study of nonreconstructed patients. *Am J Sports Med* 2007;35(7):1135–43.
31. Cameron KL, Thompson BS, Peck KY, Owens BD, Marshall SW, Svoboda SJ. Normative values for the KOOS and WOMAC in a young athletic population: history of knee ligament injury is associated with lower scores. *Am J Sports Med* 2013;41:582–9.
32. Roos EM, Bremander AB, Englund M, Lohmander LS. Change in self-reported outcomes and objective physical function over 7 years in middle-aged subjects with or at high risk of knee osteoarthritis. *Ann Rheum Dis* 2008;67(4):505–10.
33. Paradowski PT, Englund M, Lohmander S, Roos EM. The effect of patient characteristics on variability in pain and function over two years in early knee osteoarthritis. *Health Qual Life Outcomes* 2005;3:59.
34. Sackett DL, Richardson WS, Rosenberg W. Evidence-based Medicine: How to Practice and Teach EBM. New York: Churchill Livingstone; 1997.
35. Howe LD, Tilling K, Galobardes B, Lawlor DA. Loss to follow-up in cohort studies: bias in estimates of socioeconomic inequalities. *Epidemiology* 2013;24(1):1–9.
36. Winding TN, Andersen JH, Labriola M, Nohr EA. Initial non-participation and loss to follow-up in a Danish youth cohort: implications for relative risk estimates. *J Epidemiol Community Health* 2014;68(2):137–44.
37. Larsson S, Englund M, Struglics A, Lohmander LS. The association between changes in synovial fluid levels of ARGS-aggregan fragments, progression of radiographic osteoarthritis and self-reported outcomes: a cohort study. *Osteoarthr Cartil* 2012;20:388–95.
38. Oiestad BE, Holm I, Engebretsen, Risberg MA. The association between radiographic knee osteoarthritis and knee symptoms, function and quality of life 10–15 years after anterior cruciate ligament reconstruction. *Br J Sports Med* 2011;45:583–8.